

## Tagungsnummer

P53

## Thema

AG Bodengase

Neue Entwicklungen bei Methoden zur Messung und bei der Modellierung von Spurengasflüssen

## Autoren

S. Wirth<sup>1</sup>, M. Hoffmann<sup>2</sup>, H. Beßler<sup>3</sup>, C. Engels<sup>3</sup>, H. Jochheim<sup>4</sup>, M. Sommer<sup>5</sup>, J. Augustin<sup>1</sup>

<sup>1</sup>Leibniz-Zentrum für Agrarlandschaftsforschung (ZALF) e.V., Institut für Landschaftsbiogeochemie, Müncheberg; <sup>2</sup>Leibniz-Zentrum für Agrarlandschaftsforschung (ZALF) e.V., Forschungs-Informationssysteme, Müncheberg; <sup>3</sup>Humboldt University Berlin, Albrecht-Daniel-Thaer Institut, Berlin; <sup>4</sup>Leibniz-Zentrum für Agrarlandschaftsforschung (ZALF) e.V., Institut für Landschaftssystemanalyse, Müncheberg; <sup>5</sup>Leibniz-Zentrum für Agrarlandschaftsforschung (ZALF) e.V., Institut für Bodenlandschaftsforschung, Müncheberg

## Titel

Combining a root exclusion technique with continuous measurements of CO<sub>2</sub> by chambers and inside soil for a pin-point separation of ecosystem respiration in croplands

## Abstract

To better assess ecosystem C budgets of croplands and understand their potential response to climate and management changes, detailed information on the mechanisms and environmental controls driving the individual C flux components are needed. This accounts in particular for the ecosystem respiration ( $R_{eco}$ ) and its components, the autotrophic ( $R_a$ ) and heterotrophic respiration ( $R_h$ ) which vary tremendously in time and space. Therefore, we developed and tested a method to separate  $R_{eco}$  into  $R_a$  (as the sum of  $R_{a(shoot)}$  and  $R_{a(root)}$ ) and  $R_h$  in order to detect temporal and small-scale spatial dynamics within their relative contribution to overall  $R_{eco}$ . Investigations were carried out for winter wheat (*Triticum aestivum*) during the crop season 2015 at an experimental plot (CarboZALF-D) located in the hummocky ground moraine landscape of NE Germany.  $R_{eco}$  was derived from CO<sub>2</sub> flux measurements from plant stand and soil during nighttime using automatic chambers.  $R_h$  was derived from CO<sub>2</sub> efflux measurements from fallow next to the automatic chambers using CO<sub>2</sub> sampling tubes in 10 cm soil depth.  $R_{a(root)}$  was calculated as the difference between CO<sub>2</sub> efflux measurements in planted soil and  $R_h$ .  $R_{a(shoot)}$  was calculated as  $R_{eco} - R_{a(root)} - R_h$ .  $R_{eco}$  varied seasonally from <1 to 9.5 g C m<sup>-2</sup> d<sup>-1</sup>, and was higher in adult (a) and reproductive (r) than juvenile (j) stands (g C m<sup>-2</sup> d<sup>-1</sup>: j 1.2, a 4.6, r 5.3). Observed  $R_a$